


1.)

$$\lim_{x \rightarrow 0} \frac{4x^2}{e^{4x} - 4x - 1} \text{ is}$$

- (A) 0 (B) $\frac{1}{2}$ (C) 8 (D) nonexistent


2.)




Graph of f'

The graph of f' , the derivative of the function f , is shown above. Which of the following could be the graph of f ?


(A)




(B)



(C)



(D)



3.)

Let f be the function given by $f(x) = x^3 - 6x^2 - 15x$. What is the maximum value of f on the interval $[0, 6]$?


- (A) 0 (B) 5 (C) 6 (D) 8

4.)

Let f be the function defined by $f(x) = \sqrt[3]{x}$. What is the approximation for $f(10)$ found by using the line tangent to the graph of f at the point $(8, 2)$?

- (A) $\frac{11}{6}$ (B) $\frac{25}{12}$ (C) $\frac{13}{6}$ (D) $\frac{7}{3}$

5.)



Graph of f

The graph of the function f is shown in the figure above. Which of the following statements must be false?

- (A) $\lim_{x \rightarrow 2^-} f(x) = 3$
 (B) $\lim_{x \rightarrow 2^+} f(x) = \infty$
 (C) $\lim_{x \rightarrow 2} f(x) = f(2)$
 (D) $\lim_{x \rightarrow \infty} f(x) = 0$

- 6.) If $f(x) = (x^2 + 1)^3$, what is $\lim_{x \rightarrow -1} \frac{f(x) - f(-1)}{x + 1}$?
(A) -24 (B) -8 (C) 0 (D) 12

- 7.) If $y = x^2(e^x - 1)$, then $\frac{dy}{dx} =$
(A) $2xe^x$
(B) $2xe^x - 2x$
(C) $x^2e^x + 2xe^x - 2x$
(D) $x^2e^x + 2xe^x - x^2 - 2x$

- 8.) When $x = 2e$, $\lim_{h \rightarrow 0} \frac{\ln(x+h) - \ln(x)}{h}$ is
(A) $\frac{1}{2e}$ (B) 1 (C) $\ln(2e)$ (D) nonexistent

- 9.) A spherical snowball is melting in such a way that it maintains its shape. The snowball is decreasing in volume at a constant rate of 8 cubic centimeters per hour. At what rate, in centimeters per hour, is the radius of the snowball decreasing at the instant when the radius is 10 centimeters? (The volume of a sphere of radius r is $V = \frac{4}{3}\pi r^3$.)
(A) $\frac{1}{50\pi}$ (B) $\frac{3}{50\pi}$ (C) 400π (D) 3200π

10.)

Let f be the function given by $f(x) = 2 \cos x + 1$. What is the approximation for $f(1.5)$ found by using the line tangent to the graph of f at $x = \frac{\pi}{2}$?

- (A) -2 (B) 1 (C) $\pi - 2$ (D) $4 - \pi$

11.)

If $f(x) = \sin^{-1} x$, then $f'\left(\frac{\sqrt{3}}{2}\right) =$


- (A) $\frac{\pi}{6}$ (B) $\frac{\pi}{3}$ (C) $\frac{4}{7}$ (D) 2

12.)

If $f(x) = \ln x$, then $\lim_{x \rightarrow 3} \frac{f(x) - f(3)}{x - 3}$ is

- (A) $\frac{1}{3}$ (B) e^3 (C) $\ln 3$ (D) nonexistent

13.)



Graph of f

The graph of $y = f(x)$ is shown above. What is $\lim_{x \rightarrow 1} f(x)$?

- (A) 0 (B) 1 (C) 3 (D) The limit does not exist.

14.)

x	3	7
$h(x)$	7	22
$h'(x)$	5	10

Selected values of the increasing function h and its derivative h' are shown in the table above. If g is a differentiable function such that $h(g(x)) = x$ for all x , what is the value of $g'(7)$?


- (A) $-\frac{1}{10}$ (B) $\frac{1}{10}$ (C) $\frac{1}{5}$ (D) $\frac{7}{5}$

15.)

$$\lim_{x \rightarrow -7} \frac{x+7}{|x+7|} \text{ is}$$

- (A) -1 (B) 0 (C) 1 (D) nonexistent

16.)

Graph of g

The graph of the function g is shown above. If f is the function given by $f(x) = g(g(x))$, what is the value of $f'(0)$?

- (A) -2 (B) -1 (C) 2 (D) 3

17.)

Let g be a twice-differentiable, increasing function of t . If $g(0) = 20$ and $g(10) = 220$, which of the following must be true on the interval $0 < t < 10$?

- (A) $g'(t) = 0$ for some t in the interval.
 (B) $g'(t) = 20$ for some t in the interval.
 (C) $g''(t) = 0$ for some t in the interval.
 (D) $g''(t) > 0$ for all t in the interval.

18.)

Let $y = f(x)$ be a differentiable function such that $\frac{dy}{dx} = \frac{x}{y}$ and $f(8) = 2$. What is the approximation of $f(8.1)$ using the line tangent to the graph of f at $x = 8$?

- (A) 0.4 (B) 2.025 (C) 2.4 (D) 6

19.)

The number of gallons of water in a storage tank at time t , in minutes, is modeled by $w(t) = 25 - t^2$ for $0 \leq t \leq 5$. At what rate, in gallons per minute, is the amount of water in the tank changing at time $t = 3$ minutes?

- (A) 66 (B) 16 (C) -3 (D) -6

20.)

For $t \geq 0$, the velocity of a particle moving along the x -axis is given by $v(t) = t^3 - 6t^2 + 10t - 4$. At what time t does the direction of motion of the particle change from right to left?

- (A) 0.586 (B) 1.184 (C) 2.000 (D) 2.816

21.)

If $f'(x) = 3x^2 + 2x$ and $f(2) = 3$, then $f(1) =$

- (A) -10 (B) -7 (C) 10 (D) 13

22.)

What is the absolute minimum value of $y = -\cos x - \sin x$ on the closed interval $\left[0, \frac{\pi}{2}\right]$?

- (A) $-2\sqrt{2}$ (B) -2 (C) $-\sqrt{2}$ (D) -1

23.)

A particle moves along the x -axis so that at time $t > 0$ its position is given by $x(t) = 12e^{-t}\sin t$. What is the first time t at which the velocity of the particle is zero?


- (A) $\frac{\pi}{4}$ (B) $\frac{\pi}{2}$ (C) $\frac{3\pi}{4}$ (D) π

24.)

$$\lim_{x \rightarrow \infty} \frac{\ln(e^{3x} + x)}{x} =$$

- (A) 0 (B) 1 (C) 3 (D) ∞

25.)



The graph of f' , the derivative of the function f , is shown in the figure above. Which of the following statements must be true?

- I. f is continuous on the open interval (a, b) .
 II. f is decreasing on the open interval (a, b) .
 III. The graph of f is concave down on the open interval (a, b) .
- (A) I only
 (B) I and II only
 (C) I and III only
 (D) II and III only

26.)

Which of the following is a solution to the differential equation $y'' - 4y = 0$?

- (A) $y = e^{2x}$ (B) $y = 2e^x$ (C) $y = \sin(2x)$ (D) $y = \cos(2x)$

27.)

The function f has first derivative given by $f'(x) = x^4 - 6x^2 - 8x - 3$. On what intervals is the graph of f concave up?

- (A) $(2, \infty)$ only
 (B) $(0, \infty)$
 (C) $(-1, 2)$
 (D) $(-\infty, -1)$ and $(3, \infty)$

28.)

x	1	2	3	4	5
$f(x)$	9	4	0	-3	-5

The table above gives values of a function f at selected values of x . If f is twice-differentiable on the interval $1 \leq x \leq 5$, which of the following statements could be true?

- (A) f' is negative and decreasing for $1 \leq x \leq 5$.
 (B) f' is negative and increasing for $1 \leq x \leq 5$.
 (C) f' is positive and decreasing for $1 \leq x \leq 5$.
 (D) f' is positive and increasing for $1 \leq x \leq 5$.

29.)

A tire that is leaking air has an initial air pressure of 30 pounds per square inch (psi). The function $t = f(p)$ models the amount of time t , in hours, it takes for the air pressure of the tire to reach p psi. What are the units for $f'(p)$?

- (A) hours (B) psi (C) psi per hour (D) hours per psi

30.)

An isosceles right triangle with legs of length s has area $A = \frac{1}{2}s^2$. At the instant when $s = \sqrt{32}$ centimeters, the area of the triangle is increasing at a rate of 12 square centimeters per second. At what rate is the length of the hypotenuse of the triangle increasing, in centimeters per second, at that instant?

- (A) $\frac{3}{4}$ (B) 3 (C) $\sqrt{32}$ (D) 48

31.)

The graph of which of the following functions has exactly one horizontal asymptote and no vertical asymptotes?

- (A) $y = \frac{1}{x^2 + 1}$
 (B) $y = \frac{1}{x^3 + 1}$
 (C) $y = \frac{1}{e^x - 1}$
 (D) $y = \frac{1}{e^x + 1}$

32.)

The continuous function f is positive and has domain $x > 0$. If the asymptotes of the graph of f are $x = 0$ and $y = 2$, which of the following statements must be true?

- (A) $\lim_{x \rightarrow 0^+} f(x) = \infty$ and $\lim_{x \rightarrow 2} f(x) = \infty$
 (B) $\lim_{x \rightarrow 0^+} f(x) = 2$ and $\lim_{x \rightarrow \infty} f(x) = 0$
 (C) $\lim_{x \rightarrow 0^+} f(x) = \infty$ and $\lim_{x \rightarrow \infty} f(x) = 2$
 (D) $\lim_{x \rightarrow 2} f(x) = \infty$ and $\lim_{x \rightarrow \infty} f(x) = 2$



33.)

Which of the following limits are equal to -1 ?

- I. $\lim_{x \rightarrow 0^-} \frac{|x|}{x}$
 II. $\lim_{x \rightarrow 3} \frac{x^2 - 7x + 12}{3 - x}$
 III. $\lim_{x \rightarrow \infty} \frac{1 - x}{1 + x}$

- (A) I only (B) I and III only (C) II and III only (D) I, II, and III

34.)



The graphs of the functions f and g are shown in the figures above. Which of the following statements is false?

- (A) $\lim_{x \rightarrow 1} f(x) = 0$
- (B) $\lim_{x \rightarrow 2} g(x)$ does not exist.
- (C) $\lim_{x \rightarrow 1} (f(x)g(x+1))$ does not exist.
- (D) $\lim_{x \rightarrow 1} (f(x+1)g(x))$ exists.

35.)

The positive variables p and c change with respect to time t . The relationship between p and c is given by the equation $p^2 = (20 - c)^3$. At the instant when $\frac{dp}{dt} = 41$ and $c = 15$, what is the value of $\frac{dc}{dt}$?

- (A) $-\frac{82}{75}$ (B) $-\frac{2\sqrt{5}}{3}$ (C) $-\frac{3\sqrt{5}}{2}$ (D) $-\frac{82\sqrt{5}}{15}$

36.)

$$\lim_{x \rightarrow -\infty} \frac{3 + 2^x}{4 - 5^x} \text{ is}$$

- (A) $-\frac{2}{5}$ (B) 0 (C) $\frac{3}{4}$ (D) nonexistent

37.)

For time $t \geq 1$, the position of a particle moving along the x -axis is given by $p(t) = \sqrt{t} - 2$. At what time t in the interval $1 \leq t \leq 16$ is the instantaneous velocity of the particle equal to the average velocity of the particle over the interval $1 \leq t \leq 16$?

- (A) 1 (B) $\frac{121}{25}$ (C) $\frac{25}{4}$ (D) 25

38.)

If f is a differentiable function and $y = \sin(f(x^2))$, what is $\frac{dy}{dx}$ when $x = 3$?

- (A) $\cos(f'(9))$
 (B) $6 \cos(f(9))$
 (C) $f'(9) \cos(f(9))$
 (D) $6f'(9) \cos(f(9))$

39.)

The number of insects in a certain population at time t days is modeled by the function P with first derivative $P'(t) = 0.3t^2 + 12t + 210$. At time $t = 0$, the number of insects in the population is 40. Which of the following statements are true?

- I. At time $t = 10$, the number of insects in the population is 2840.
 II. At time $t = 10$, the number of insects in the population is increasing at a rate of 360 insects per day.
 III. At time $t = 10$, the rate of change of the number of insects in the population is increasing at a rate of 18 insects per day per day.

- (A) I only (B) II only (C) III only (D) I, II, and III

40.)

If $f(x) = \cos^2(3x - 5)$, then $f'(x) =$

- (A) $6 \cos(3x - 5)$
 (B) $-3 \sin^2(3x - 5)$
 (C) $-2 \sin(3x - 5) \cos(3x - 5)$
 (D) $-6 \sin(3x - 5) \cos(3x - 5)$

41.)

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	-0.1054	-0.0101	-0.001	0.001	0.0099	0.0953

The function f is continuous and increasing for $x > -1$. The table above gives values of f at selected values of x . Of the following, which is the best approximation for $\lim_{x \rightarrow 0} e^{-2f(x)}$?

- (A) -2
- (B) 0
- (C) 1
- (D) The limit does not exist.

42.)

x	10	11	12	13	14
$f(x)$	5	2	3	6	5

The table above gives values of the continuous function f at selected values of x . If f has exactly two critical points on the open interval $(10, 14)$, which of the following must be true?

- (A) $f(x) > 0$ for all x in the open interval $(10, 14)$.
- (B) $f'(x)$ exists for all x in the open interval $(10, 14)$.
- (C) $f'(x) < 0$ for all x in the open interval $(10, 11)$.
- (D) $f'(12) \neq 0$

43.)

Let f be the function with $f(0) = \frac{1}{\pi^2}$, $f(2) = \frac{1}{\pi^2}$, and derivative given by $f'(x) = (x+1)\cos(\pi x)$. How many values of x in the open interval $(0, 2)$ satisfy the conclusion of the Mean Value Theorem for the function f on the closed interval $[0, 2]$?

- (A) None
- (B) One
- (C) Two
- (D) More than two

44.)

Let f be a twice-differentiable function for all real numbers x . Which of the following additional properties guarantees that f has a relative minimum at $x = c$?

- (A) $f'(c) = 0$
- (B) $f'(c) = 0$ and $f''(c) < 0$
- (C) $f'(c) = 0$ and $f''(c) > 0$
- (D) $f'(x) > 0$ for $x < c$ and $f'(x) < 0$ for $x > c$

45.)

The rate at which water leaks from a tank, in gallons per hour, is modeled by R , a differentiable function of the number of hours after the leak is discovered. Which of the following is the best interpretation of $R'(3)$?

- (A) The amount of water, in gallons, that has leaked out of the tank during the first three hours after the leak is discovered
- (B) The amount of change, in gallons per hour, in the rate at which water is leaking during the three hours after the leak is discovered
- (C) The rate at which water leaks from the tank, in gallons per hour, three hours after the leak is discovered
- (D) The rate of change of the rate at which water leaks from the tank, in gallons per hour per hour, three hours after the leak is discovered

46.)

Let f be a twice-differentiable function such that $f''(x) < 0$ for all x . The graph of $y = S(x)$ is the secant line passing through the points $(3, f(3))$ and $(5, f(5))$. The graph of $y = T(x)$ is the line tangent to the graph of f at $x = 4$. Which of the following is true?

- (A) $f(4.2) < S(4.2) < T(4.2)$
- (B) $f(4.2) < T(4.2) < S(4.2)$
- (C) $S(4.2) < f(4.2) < T(4.2)$
- (D) $T(4.2) < f(4.2) < S(4.2)$

47.)

If $f(x) = 3x^2 + 2x$, then $f'(x) =$

- (A) $\lim_{h \rightarrow 0} \frac{(3x^2 + 2x + h) - (3x^2 + 2x)}{h}$
- (B) $\lim_{x \rightarrow 0} \frac{(3x^2 + 2x + h) - (3x^2 + 2x)}{h}$
- (C) $\lim_{h \rightarrow 0} \frac{(3(x+h)^2 + 2(x+h)) - (3x^2 + 2x)}{h}$
- (D) $\lim_{x \rightarrow 0} \frac{(3(x+h)^2 + 2(x+h)) - (3x^2 + 2x)}{h}$

48.) The function f is increasing on the interval $[1, 3]$ and nowhere else. The first derivative of f , f' , is continuous for all real numbers. Which of the following could be a table of values for $f'(x)$?

(A)

x	$f'(x)$
0	-1
1	0
2	2
3	0
4	-2

(B)

x	$f'(x)$
0	-1
1	1
2	2
3	1
4	-2

(C)

x	$f'(x)$
0	1
1	0
2	1
3	2
4	0

(D)

x	$f'(x)$
0	1
1	0
2	2
3	0
4	-2

49.) On a certain day, the total number of pieces of candy produced by a factory since it opened is modeled by C , a differentiable function of the number of hours since the factory opened. Which of the following is the best interpretation of $C'(3) = 500$?

- (A) The factory produces 500 pieces of candy during its 3rd hour of operation.
- (B) The factory produces 500 pieces of candy in the first 3 hours after it opens.
- (C) The factory is producing candy at a rate of 500 pieces per hour, 3 hours after it opens.
- (D) The rate at which the factory is producing candy is increasing at a rate of 500 pieces per hour per hour, 3 hours after it opens.

50.)

$$f(x) = \begin{cases} x^2 \sin(\pi x) & \text{for } x < 2 \\ x^2 + cx - 18 & \text{for } x \geq 2 \end{cases}$$

Let f be the function defined above, where c is a constant. For what value of c , if any, is f continuous at $x = 2$?

- (A) 2
- (B) 7
- (C) 9
- (D) $4\pi - 4$
- (E) There is no such value of c .